



Discharge and Transition Planning – NICU and Pediatrics Bibliography

- Ahmann, E., Abraham, M. R., & Johnson, B. H. (2003). *Changing the concept of families as visitors: Supporting family presence and participation*. Bethesda, MD: Institute for Patient- and Family-Centered Care. Available from <http://www.ipfcc.org/resources/pinwheel/index.html>
- Ahmann, E., Webster, P. D., & Johnson, B. H. (2000). *Creating and expanding patient and family resource centers*. Bethesda, MD: Institute for Patient- and Family-Centered Care. Available from <http://www.ipfcc.org/resources/pinwheel/index.html>
- American Academy of Pediatrics. (1998). Hospital discharge of the high-risk neonate—Proposed guidelines (RE9812). *Pediatrics*, 102(2), 411-417.
- American Heart Association. (2000). Guidelines 2000 for cardiopulmonary resuscitation and emergency cardiovascular care: International consensus on science. *Circulation*, 102 (Suppl. I): I-1-I-384. Available from <http://circ.ahajournals.org/content/vol102/issue90001/index.shtml>
- Baroni, M. A., Tuthill, P., Feenan, L., & Schroeder, M. (1994). Technology-dependent infants and young children: A retrospective care analysis of service coordination across state lines. *Infants and Young Children*, 7(1), 69-78.
- Brazy, J. E., Anderson, B. M., Becker, P. T., & Becker, M. (2001). How parents of premature infants gather information and obtain support. *Neonatal Network*, 20(2), 41-48.
- Broedsgaard, A., & Wagner, L. (2005). How to facilitate parents and their premature infant for the transition home. *International Nursing Review*, 52, 196-203.
- Browne, J. V., Langlois, A., Ross, E. S., & Smith-Sharp, S. (2001). Beginnings: An interim individualized family service plan for use in the intensive care nursery. *Infants and Young Children*, 14(2), 19-32.
- Brun, D. A., & McCollum, J. A. (2002). Partnerships between mothers and professionals in the NICU: Caregiving, information exchange and relationships. *Neonatal Network*, 21(7), 15-23.
- Conner, J. M., & Nelson, E. C. (1999). Neonatal intensive care: Satisfaction measured from a parent's perspective. *Pediatrics*, 103(1 Su), 336-349.
- Cox, M. (1996). Discharge planning for the high-risk infant. In E. Ahmann (Ed.). *Home care for the high-risk infant: A family-centered approach* (pp. 45-51). Gaithersburg, MD: Aspen.
- Dokken, D. L., & Sydnor-Greenberg, N. (1998). Helping families mobilize their personal resources. *Pediatric Nursing*, 24(1), 66-69.
- Forsythe, P. L., Maher, R., Kirchick, C., & Bieda, A. (2007). SAFE discharge for infants with high-risk home environments. *Advances in Neonatal Care*, 7(2), 69-75.
- Gaal, B. J., Blatz, S., Dix, J., & Jennings, B. (2008). Discharge planning utilizing the discharge train: Improved communication with families. *Advances in Neonatal Care*, 8(1), 42-55.
- Gannon, B. (2000). Caring one day at a time. *Neonatal Network*, 19(5), 25-32.
- Griffin, T., & Abraham, M. R. (2006). Transition to home from the newborn intensive care unit: Applying the principles of family-centered care to the discharge process. *Journal of Perinatal and Neonatal Nursing*, 20(3), 243-249.

- Hadden, D. S. (2000). The long anticipated day: Strategies for success when a premature infant comes home from the neonatal intensive care unit. *Young Exceptional Children*, 3(2), 21-26.
- Hostler, S. L. (1999). Pediatric family-centered rehabilitation.** *Journal of Head Trauma Rehabilitation*, 14, 384-393.
- Hussey-Gardner, B., McNinch, A., Anastasi, J. M., & Miller, M. (2002). Early intervention best practice: Collaboration among an NICU, an early intervention program and an NICU follow-up program. *Neonatal Network*, 21(3), 15-22.
- Institute for Patient- and Family-Centered Care. (2003). *Partnerships with families in newborn intensive care...Enhancing quality and safety* [Videotape]. Available from <http://www.ipfcc.org/resources/videos/index.html>
- Johnson, B. H., & Eichner, J. M. (2003). Family-centered care and the pediatrician's role. *Pediatrics*, 112(3), 691-696.
- Johnson, B. H., & Hanson, J. L. (Executive Producers). (1996). *Newborn intensive care: Changing practice, changing attitudes*. [Videotape]. Available from <http://www.ipfcc.org/resources/videos/index.html>
- Johnston, A. M., Bullock, C. E., Graham, J. E., Reilly, M. C., Rocha, C., Hoopes, R. D., Jr., et al. (2006). Implementation of potentially better practices for family-centered care: The family-centered care map. *Pediatrics*, 118(Suppl. 2), s108-s114.
- Lewis, M., & Noyes, J. (2007). Discharge management for children with complex needs. *Paediatric Nursing*, 19(4), 26-30.
- Mancini, A., & While, A. (2001). Discharge planning from a neonatal unit: An exploratory study of parents' views. *Journal of Neonatal Nursing*, 7(2) 59-62.
- Noyes, J., & Lewis, M. (2005). *Care pathway for the discharge and support of children requiring long-term ventilation in the community*. London, UK: Department of Health. Available from http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_4115098
- Saunders, R. P., Abraham, M. R., Crosby, M. J., Thomas, K., & Edwards, W. H. (2003). Evaluation and development of potentially better practices for improving family-centered care units. *Pediatrics*, 111(4 pt. 2), e437-e449.
- Webster, P. D., & Johnson, B. H. (1999). *Collaborative design planning*. Bethesda, MD: Institute for Patient- and Family-Centered Care. Available from <http://www.ipfcc.org/resources/pinwheel/index.html>
- Webster, P. D., & Johnson, B. H. (2000). *Developing and sustaining a patient and family advisory council*. Bethesda, MD: Institute for Patient- and Family-Centered Care. Available from <http://www.ipfcc.org/resources/pinwheel/index.html>